



March 2008 (updated May 2015)

ELDER CARE

A Resource for Interprofessional Providers



A program of the Hartford
Geriatric Nursing Initiative

Depression in Elders

M. Jane Mohler, MPH, PHD, College of Medicine, University of Arizona

Depression is commonly described as feeling sad, blue, unhappy, miserable, or down in the dumps. While many of us feel this way for short periods, true clinical depressive disorders are syndromes characterized by the impairment of mood regulation. The most common diagnoses include major depression and dysthymia, a disorder characterized by chronic low mood. Prevalent among older adults, depression is associated with a 1.5–3 times higher incidence of medical morbidity, and the lifetime risk of suicide is reported to be as high as 15%. Depression negatively affects functioning and quality of life, contributes to excess morbidity and mortality, and places extra stress on caregivers and the health care system. Of the estimated 6 million persons over age 65 with depression, only 10% receive treatment. In addition, less than half of hospitalized patients with depression are referred to a psychiatrist, and less than 20% of these are prescribed antidepressant medication.

Epidemiology

Depression is the most common geriatric psychiatric disorder, and can manifest as either minor or major depression. Eight to fifteen percent of the general population over 65 years of age has symptoms severe enough to meet diagnostic criteria for a depressive disorder, and the prevalence of major depressive disorder (MDD) is estimated to be 2%. In the hospitalized population, however, 25–40% manifest minor depressive symptoms. In assisted living and skilled nursing facilities an estimated 30% display mild depressive symptoms and an additional 12% of patients have MDD.

Risk Factors

Major risk factors for depression include the following: female gender, bereavement, stressful life events, social isolation, chronic pain, a past

history of depression, fear of death, chronic disease, substance abuse, including alcohol, and being unmarried, widowed or cohabitating.

Signs and Symptoms


Depression in older patients can be difficult to diagnose, as signs and symptoms differ from younger adults, and may not be in accord with DSM-IV or ICD-10 criteria. Additionally, medical illnesses can confound the symptoms of depression. Older adults may not show or express sadness, their mood can be chronically irritable, and depressed elders can lose their ability to respond to positive external events. Somatic complaints and hypochondriasis are more frequent, and vegetative signs such as anorexia and weight loss may initiate concerns about underlying malignancy. About 10% of depressed elders may display psychotic symptoms. Between 38–58% of aging adults suffering from MDD also have anxiety disorder, which often presents as tension, unrest, feelings of insecurity or fear, irritability, and intense worry rather than as autonomic symptoms.

Screening and Diagnosis

The US Preventive Task Force recommends screening adults for depression only where “there are systems in place to assure accurate diagnosis, effective treatment and careful follow-up care.” A quick, easy, two sentence screening tool is offered under Provider Tips. A positive result should prompt further evaluation and diagnosis. Remember that medical illness can sometimes present as depression. Recommended lab tests to rule out other causes include thyroid, liver and kidney function tests, serum calcium, and B-12 levels (or homocysteine and methylmalonic acid levels). Organic cognitive disorders must also be

Signs and Symptoms of Depression in Older Adults

- Lack of appetite
- Weight loss
- Fatigue
- Problems with concentration
- Stopping normal activities
- Guilt
- Melancholia
- Suicidal Ideation
- Feeling helpless
- Feeling hopeless



Suicide Risks in Those >50 Years

- Poor Health
- Family conflict
- Money worries
- Male
- White
- Veterans
- >84 yrs have twice the risk

TIPS FOR DIAGNOSING DEPRESSION IN OLDER ADULTS

Asking these two questions may be as effective as using longer screening tools:

- Over the past 2 weeks, have you ever felt down, depressed, or hopeless?
- Over the past 2 weeks, have you felt little interest or pleasure in doing things?

A positive response to either question is a very sensitive indicator of depression, but needs further validation with a more specific diagnostic interview.

ELDER CARE

Continued from front page

considered when making a diagnosis of depression. It may be difficult to differentiate depression from dementia, and they may co-exist. Depression can precede, accompany or masquerade as dementia, and treatment of the depression will often improve cognitive functioning. Neuropsychiatric evaluation can help to tease out depression from cognitive deficit. Acute neuropsychiatry or geriatric psychiatry referral should be sought when a patient is suicidal or homicidal, has delusions or hallucinations, or is disabled by vegetative depression.

Treatment

Antidepressant medications are usually the first line of treatment for depressed older adults. Symptom improvement is well documented with serotonin reuptake inhibitors (SSRIs), tricyclic antidepressants (TCAs), and the newer drugs such as mirtazapine, nefazodone, and venlafaxine. Unfortunately, antidepressant medication often takes up to 3 months to demonstrate effect. If, after an 8-12 week therapeutic trial of one drug shows no effect, alternative medications should be prescribed. Once improvement is obtained, a minimum of 6 months of treatment is recommended. At that point a slow and carefully monitored weaning regimen may be attempted. Older patients, especially those with a history of previous depressive episodes, may require a longer duration of treatment. Occasionally combination therapy may be needed, but requires careful monitoring. A geropsychiatry consult can help in these situations. A future provider fact sheet will review the currently available medications with regard to mechanism of action, side effects, and individual indications for use. Uses for electroconvulsive therapy will also be addressed.

Psychotherapy can also benefit patients, and small additional therapeutic gains are seen when provided in combination with antidepressants. The evidence base is unclear with regard to the ideal type of psychotherapy, but those aimed at relieving depressive ideation, and efforts focused

on instrumental activities of daily living, can improve geriatric depression.

Older adults, however, may refuse therapy, as many in this present generation consider therapy a sign of weakness. Patients with neurocognitive disorder, psychomotor retardation or sensory impairment may not be suitable for psychotherapy.

A meta-analysis of MDD trials provides evidence that omega-3 supplementation is safe and reduces symptoms of depression in combination with routine care. In addition, B12 insufficiency has been linked to depression in several small studies. B vitamin supplements are not yet recommended as standard of care, but, as with omega-3s, are easy to prescribe without fear of adverse effects.

Special Considerations

Bereavement

Those going through uncomplicated bereavement are likely to experience a lack of energy and concentration, crying spells, and decreased appetite and insomnia. Most will need no formal intervention. Occasionally, such depression may deepen, resulting in overwhelming feelings of sadness, sometimes to the point of suicidal ideation. In this case, it can be helpful to talk with clergy or spiritual healers, or with a social worker, grief counselor, or therapist. Support groups can also be helpful. Antidepressants and counseling have been found to be effective in combination in grief.

Suicide

Nearly 25% of elders suffering from MDD will remit, either spontaneously or after treatment. Another 25% will not respond to any kind of intervention and will continue to manifest severe symptoms. The other 50% will have partial remission, or intermittent recurrence. MDD accounts for 65% of cases of elderly suicide. Screening for suicidal ideation in a depressed older adult is paramount. An acute life threatening illness (e.g., MI, stroke, or cancer diagnosis) may trigger suicidal plans. Don't be afraid to ask.

References and Resources

- Chapman DP, Perry GS. Depression as a major component of public health for older adults. *Prev Chronic Dis*. 2008;5 (1):A22.
- Fountoulakis KN, O'Hara R, Iacovides A, et al. Unipolar late-onset depression: A comprehensive review. *Ann Gen Hosp Psychiatry*. 2003;2(1):11. 10.1186/1475-2832-2-11.
- Schneider L. *Diagnosis and Treatment of Depression in Late Life: Results of the NIH Consensus Development Conference*. 1st ed. Washington, DC: American Psychiatric Press; 1994.
- Screening for Depression, Topic Page. May 2002. U.S. Preventive Services Task Force. Agency for Healthcare Research and Quality, Rockville, MD. <http://www.ahrq.gov/clinic/uspstf/uspstfdepr.htm>

Interprofessional care improves the outcomes of older adults with complex health problems

Editors: Mindy Fain, MD; Jane Mohler, NP-c, MPH, PhD; and Barry D. Weiss, MD
Interprofessional Associate Editors: Tracy Carroll, PT, CHT, MPH; David Coon, PhD; Jeannie Lee, PharmD, BCPS;
Lisa O'Neill, MPH; Floribella Redondo; Laura Vitkus, BA

The University of Arizona, PO Box 245069, Tucson, AZ 85724-5069 | (520) 626-5800 | <http://aging.medicine.arizona.edu>

Supported by: Donald W. Reynolds Foundation, Arizona Geriatric Education Center and Arizona Center on Aging

This project was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number UB4HP19047, Arizona Geriatric Education Center. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.