



May 2015

ELDER CARE

A Resource for Interprofessional Providers



A program of the Hartford
Geriatric Nursing Initiative

Disease Screening in Older Adults: When to Stop

Charles P Mouton, MD, MS, School of Medicine, Meharry Medical College

National organizations, such as the US Preventive Services Task Force (USPSTF), have published recommendations for screening adults for a variety of diseases, based on a consideration of whether screening will lead to improved outcomes. Many of the recommendations are aimed at diseases that are common in older adults. While the USPSTF and other organizations do an excellent job of elucidating when and how to perform screening, less attention is given to when screening should stop. Little evidence and few recommendations are available on which to base decisions about screening and other preventive care in older adults, particularly for those over age 70. The most current information is shown in Table 1.

When choosing to screen for a particular disease, several factors need to be considered. First, the disease must be sufficiently common and serious to present a problem to individuals and to society. Second, there must be a latent or asymptomatic stage during which treatment will be of more benefit than after symptoms occur. Third, there must be a sufficiently accurate, safe, and cost-effective screening test that is acceptable to patients.

When applying screening tests to older adults, three other factors must also be considered: life expectancy, the ability to undergo the screening test and any ensuing treatments, and, finally, personal preference.

Life Expectancy

Average life expectancy in the US is now about 78 years. Individuals age at different rates, however, leading to considerable variability in survival, even at advanced ages. Vigorous older adults may benefit from many of the standard available screening tests, whereas those with multiple co-morbidities might not live long enough to profit from screening. It is important to note that the average life expectancy of nursing home residents is less than 3 years. Therefore, screening in nursing home residents should be individualized rather than routine.

Because life expectancy is not solely a function of chronologic age, many experts have suggested calculating an estimated life expectancy as a function of health, or “physiologic age.” One easy-to-use method adjusts chronologic to physiologic age by using patients’ ratings of their global health status. Patients are asked: “How is your health – excellent, good, fair, or poor?” Depending on the response, physiologic age can be determined from Table 2 and the results used to estimate life expectancy (Table 3). Other approaches expectancy have use gait speed or self-reported functional status.

Ability to Participate in Screening and Treatments

Some screening tests represent a challenge for older adults with functional limitations. For example, DEXA testing for osteoporosis may be difficult for individuals with mobility issues or those who cannot lie flat. Patients with dementia may be uncooperative and unwilling to participate in the testing procedure. All aspects of patients’ health should be considered prior to ordering a screening test.

Similarly, if a screening test detects a disease whose treatment presents an unacceptable burden to a patient, screening may not be warranted. Information about potential treatments should be discussed beforehand.

Personal Preferences

Personal preferences are an essential part of decisions about screening and most patients indicate they would continue screening throughout their life, often in spite of a clinician’s recommendation against screening. The American Geriatrics Society recommends that older patients should have access to screening tests, even if they do not plan to pursue treatment of a condition that is found. They can still incorporate the test results into planning for their remaining life course, and they should not be denied access to desired testing solely because the test results will not change management.

TIPS ABOUT SCREENING TESTS FOR OLDER ADULTS

- Avoid screening tests for people whose life expectancy is so short that they are likely to die before benefiting from treating the condition that would be detected by screening.
- Estimate life expectancy from physiologic age, not chronologic age (Tables 2 and 3).
- Order screening tests for nursing-home patients on an individual basis, as their average life expectancy is <3 yrs.

ELDER CARE

Continued from front page

Table 1: Recommendations for Screening and Stopping Screening in Older Adults (USPSTF = US Preventive Services Task Force; LE = life expectancy)

CONDITION	RECOMMENDATION	WHEN TO STOP	MODIFYING FACTORS
Dementia	Varying recommendations		Assess whenever there is clinical suspicion
Aortic aneurysm	Once in smokers of ≥ 100 cigs	After age 75	Recommendation refers only to abdominal aneurysm
Breast cancer	Every 2 years	After age 75	If LE >5 yr and can tolerate treatment, continue screening
Cervical cancer	Every 1-3 years	After age 65	Continue screening if cervix present and multiple sex partners
Colorectal cancer	Every 10 years	After age 85	Individualized in people between 76 - 85 yrs
Depression	Varying Recommendations		Screen whenever there is clinical suspicion
Diabetes	Annually		Continue screening indefinitely in patients with hypertension
Hearing	No screening per USPSTF		Other organizations recommend screening every 3-5 years
Hepatitis B	Periodically in high-risk groups	If risk factors no longer present	No screening if immunized against hepatitis B
Hepatitis C	Once		Only screen if born between 1949-1965
Hypertension	Every 1-2 years		If LE <1 yr, consider stopping
Lipid disorders	Every 1-5 years	After age 65	If LE >5 yr and treatment likely to benefit, continue screening
Lung cancer	Annual if ≥ 30 pack-yrs smoking	Age 80 or ≥ 15 yrs no smoking	Others suggest stopping at age 75
Obesity	Routine at office visits		Consider not checking If LE <20 yrs
Oral cancer	No specific recommendations		Some organizations recommend screening if high risk
Osteoporosis	Once (age 65 women, 70 men)		
Ovarian cancer	No screening		
Pancreas cancer	No screening		
Periph artery dz	No specific recommendations		No testing after age 65
Prostate cancer	No screening		Some recommend individualized screening
Sexually transmitted infections	No specific recommendations		Consider screening if new or multiple sex partners
Thyroid disease	No specific recommendations		
Vision	No screening per USPSTF		Other organizations recommend screening every 1-2 years
Vitamin D	No specific recommendations		Consider not checking if LE < 5 years

Table 2: Estimating Physiologic Age from Chronological Age and Self-Reported Health Status

Age In Yrs	Physiologic Age (Yrs)							
	Self Reported Health Excellent		Self Reported Health Good		Self Reported Health Fair		Self Reported Health Poor	
	M	F	M	F	M	F	M	F
65	58	60	64	64	68	66	73	72
70	62	65	69	69	73	71	78	77
75	67	70	74	74	78	76	83	82
80	72	75	79	79	83	81	85+	85+

M = Male, F = Female

Source: http://symptomresearch.nih.gov/chapter_14/Part_3/sec4/chspt3s4pg1.htm

Table 3: Life Expectancy for Men and Women in the US Based Solely on Chronological Age

Age In Yrs	Life Expectancy in Years (Percentile)					
	Male			Female		
	25 th	50 th	75 th	25 th	50 th	75 th
70	6.7	12.4	18.0	9.5	15.7	21.3
75	4.9	9.3	14.2	6.8	11.9	17.0
80	3.3	6.7	10.8	4.6	8.6	13.0
85	2.2	4.7	7.9	2.9	5.9	9.6
90	1.5	3.2	5.8	1.8	3.9	6.8
95	1.0	2.3	4.3	1.1	2.7	4.8

Data in this table are population statistics that have not been adjusted for health status (physiologic age). Use the information in Table 3 to determine an individual's physiologic age, and then use the age in this table to estimate life expectancy. Source: JAMA; 285:2751.

References and Resources

US Preventive Services Task Force Recommendations: <http://www.uspreventiveservicestaskforce.org/BrowseRec/Index/browse-recommendations>

Interprofessional care improves the outcomes of older adults with complex health problems

Editors: Mindy Fain, MD; Jane Mohler, NP-c, MPH, PhD; and Barry D. Weiss, MD

Interprofessional Associate Editors: Tracy Carroll, PT, CHT, MPH; David Coon, PhD; Jeannie Lee, PharmD, BCPS; Lisa O'Neill, MPH; Floribella Redondo; Laura Vitkus, BA

The University of Arizona, PO Box 245069, Tucson, AZ 85724-5069 | (520) 626-5800 | <http://aging.medicine.arizona.edu>

Supported by: Donald W. Reynolds Foundation, Arizona Geriatric Education Center and Arizona Center on Aging

This project was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number UB4HP19047, Arizona Geriatric Education Center. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.